


Simon, E. 1900. Descriptions d'arachnides nouveaux de la famille des Attidae. Ann. Soc. ent. Belg. 44: 397.

Hispo frenata (Simon, 1900)

Astænorchestes (1) **frenatus** sp. nov. — ♂. Long. 5 mill. — Cephalothorax niger, parte cephalica antice, thoracica in medio dilutioribus et rufescentibus, pilis albis, ad marginem frontalem et utrinque sub oculis lineam exilem, in parte thoracica lineas abbreviatas et divaricatas designantibus, ornatus, parte cephalica impressionibus latis, leviter coriaceo-striolatis, munita. Pili oculorum pauci albidii. Clypeus glaber. Abdomen angustum (valde detritum), nigrum, vittis transversis albo-pilosis ornatum. Chelæ nigro-cupreæ, nitidæ, antice deplanatæ sed costa obliqua sectæ. Sternum fusco-rufescens. Pedes quatuor antici nigricantes, tarsi 1ⁱ paris, metatarsis tarsisque 2ⁱ paris luteis. Pedes postici femoribus patellisque nigricantibus, tibiis fuscis, reliquis articulis luteis sed metatarsis apice minute fuscis. Pedes-maxillares nigricantes, femore ad apicem albo-piloso; tibia patella brevior, apophysi bifida: ramulo superiore sat longo recto gracili sed obtuso, inferiore brevi et uncatu, extus armata; tarso late ovato; bulbo maximo, ad basin valde convexo et in declivitate anteriore stylo, circulum formante, ornato. — Madagascar: Imerina (Camboué).



E. *Astænorchestes frenatus* E. Sim. Chélicère ♂ en avant.

F. *Idem*. P.-M. ♂ de profil.

[Simon, E. 1901. Histoire naturelle des araignées. Paris, 2: 450].

(1) **Astænorchestes** nov. gen. — Ab *Hispone* differt quadrangulo oculorum. haud longiore quam antice latiore, sternoque antice vix attenuato, late truncato haud depresso. — Typus *A. frenatus* E. S.